


Saving Women's Lives With Family Planning

Fewer Women Die When Pregnancies are Planned

Family planning saves women's lives in two ways. First, family planning helps women avoid unwanted pregnancies. Each year, 687,000 maternal deaths are averted with family planning versus the 533,000 maternal deaths that occur.¹ As shown in the graph,² the more couples practice family planning, the fewer women die from pregnancy-related causes.^{3,4,5} Women who are not pregnant do not die from pregnancy-related causes.

Second, family planning helps couples plan the size of their family and avoid high-risk pregnancies. Women who become pregnant when very young or very old are at a higher risk of a pregnancy-related death. Women who become pregnant soon after their last birth or who have many children also have an increased danger of maternal death. Fewer deaths will occur among pregnant women if more couples practice family planning.



Too Young, Too Old, Too Close, Too Many

- When a woman becomes pregnant before age 18 she is more likely than an older woman to develop hypertension, anemia and prolonged or obstructed labor⁶ and to die as a result of pregnancy.⁷
- For pregnancies among women in their 40s, the risk of death can be as much as seven times higher than among women in their 20s.⁷
- When pregnancy occurs less than six months from the last birth, a woman is more than twice as likely to die in childbirth than a woman who becomes pregnant between 18 to 23 months later.⁸
- Five or more pregnancies significantly increase the risk of dying of pregnancy-related causes.^{7,9,10}

Saving the Lives of Indonesia's Women

It is estimated that for every 100,000 babies born, 240 women die in Indonesia.³ This estimate indicates that progress is being made but there is more work to be done. Contraceptive use is fairly high in Indonesia with 57% of married women currently using a modern method. However, only 11% of married women report using a highly effective, long-term method of contraception despite over 50% of married women reporting not wanting anymore children.⁴ Increasing women's access to and knowledge of long-term methods would result in fewer unintended pregnancies and fewer maternal deaths each year.

What Can Be Done

- ◆ Make family planning services and supplies available for highly effective, long-term contraceptive methods in the Indonesian health system.
- ◆ Ensure that Indonesian women and couples are able to plan their families and avoid unwanted and high risk pregnancies.

Sources

- ¹Personal communication with Amy Tsui, Professor, Johns Hopkins School of Public Health and Director, Bill and Melinda Gates Institute for Population and Reproductive Health. Calculations based on Liu L, et al. Three methods of estimating births averted nationally by contraception. *Population Studies*, 2008 July; 62(2): 191-210.
- ²Calculations for maternal mortality rate by Carie Muntifering, Doctoral Candidate, Johns Hopkins School of Public Health. Calculations based on: World Health Organization. *Trends in Maternal Mortality:1990 to 2008. Estimates Developed by WHO, UNICEF, UNFPA and the World Bank*, Geneva: WHO, 2007 and *World Population Prospects: The 2008 Revision Population Database*. United Nations, 2008. <http://esa.un.org/UNPP/>, accessed 13 Dec 2010.
- ³World Health Organization. *Trends in Maternal Mortality:1990 to 2008. Estimates Developed by WHO, UNICEF, UNFPA and the World Bank*, Geneva: WHO, 2007.
- ⁴Macro International Inc. MEASURE DHS STAT compiler. <http://www.measuredhs.com>, accessed 07 Dec 2010.
- ⁵World Population Prospects: The 2008 Revision Population Database. United Nations, 2008. <http://esa.un.org/UNPP/>, accessed 15 Sept 2010.
- ⁶World Health Organization. *Report of a WHO Technical Consultation on Birth Spacing, 13–15 June 2005: Geneva, Switzerland*. Geneva: Department of Making Pregnancy Safer, Department of Reproductive Health and Research, World Health Organization, 2006.
- ⁷DaVanzo J. et al. *Pregnancy Outcomes, and Maternal Morbidity and Mortality in Matlab, Bangladesh*. RAND, Labor and Population working paper series, 2004.
- ⁸Conde-Agudelo A and Belizan JM. *Maternal morbidity and mortality associated with interpregnancy interval: cross sectional study*. *British Medical Journal*, Nov 2000; 321: 1255-1259.
- ⁹Chen LC, et al. *Maternal mortality in rural Bangladesh*. *Studies in Family Planning*, 1974, 5(11), 334-341.
- ¹⁰Koenig MA, et al. *Maternal mortality in Matlab, Bangladesh: 1976-85*. *Studies in Family Planning*, 1988,19(2), 69-80.

For more information contact:

Jennifer Carlin
Advance Family Planning Program Administrator
The Bill and Melinda Gates Institute for Population and Reproductive Health
Johns Hopkins Bloomberg School of Public Health
615 N. Wolfe Street, Room W4503
Baltimore, MD 21205
Phone: 410-502-8715
Fax: 410-955-0792
jcarlin@jhsph.edu



*Bill and Melinda Gates Institute for
Population and Reproductive Health*

